

APPENDIX 1 - Malaysia

Table A1.1. Carbon debt calculation of Malaysia case

Carbon debt due to conversion of lowland tropical rainforest		estimates					references		
Aboveground carbon stock loss		197.5	Mg C ha ⁻¹	268	254.5	201.25	157	178	126
-19% forest products		37.5	Mg C ha ⁻¹						(IPCC 2006)
subtotal		159.9	Mg C ha⁻¹						
Belowground carbon stock loss									
biomass		47.8	Mg C ha ⁻¹	13%	21%	26%	23.50%	37%	(Houghton and Hackler 2001)
soil		18.2	Mg C ha ⁻¹	47.95 [Mg C ha-1]					(Niiyama et al. 2010)
subtotal		66.0	Mg C ha⁻¹						(IPCC 2006)
Carbon stocked in oil palm plantation		35.8	Mg C ha⁻¹	36	35.5	31.5	40		(Germer and Sauerborn 2008, Murdiyarso et al. 2010, Pereira de Souza et al. 2010)
Total carbon debt		190.2	Mg C ha⁻¹						
		698.8	Mg CO₂ ha⁻¹						

LITERATURE CITED

- Germer, J. and J. Sauerborn. 2008. Estimation of the impact of oil palm plantation establishment on greenhouse gas balance. *Environment, Development and Sustainability* 10:697-716.
- Hoshizahi, K., K. Niiyama, K. Kimura, T. Yamashita, Y. Bekku, T. Okuda, E. S. Quah, and N. Supardi. 2004. Temporal and spatial variation of forest biomass in relation to stand dynamics in a mature, lowland tropical rainforest, Malaysia. *Ecological Research* 19:357-363.
- Houghton, R. A. and J. L. Hackler. 2001. *Carbon flux to the atmosphere from land-use changes: 1850-1990*. No. 5054, U.S. Department of Energy.
- Imai, N., H. Samejima, A. Langner, R. C. Ong, S. Kita, J. Titin, A. Y. C. Chung, P. Lagan, Y. F. Lee, and K. Kitayama. 2009. Co-benefits of sustainable forest management in biodiversity conservation and carbon sequestration. *Plos One* 4:e8267.
- IPCC. 2006. *Guidelines for National Greenhouse Gas Inventories*
- Miettinen, J. and S. C. Liew. 2009. Estimation of biomass distribution in Peninsular Malaysia and in the islands of Sumatra, Java and Borneo based on multi-resolution remote sensing land cover analysis. *Mitigation and Adaptation Strategies for Global Change* 14:357-373.
- Murdiyarso, D., K. Hergoualc'h, and L. V. Verchot. 2010. Opportunities for reducing greenhouse gas emissions in tropical peatlands. *Proceedings of the National Academy of Sciences of the United States of America* 107:19655-19660.
- Niiyama, K., T. Kajimoto, Y. Masuura, T. Yamashita, N. Matsuo, Y. Yashiro, A. Ripin, A. R. Kassim, and N. S. Noor. 2010. Estimation of root biomass based on excavation of individual root systems in a primary dipterocarp forest in Pasoh Forest Reserve, Peninsular Malaysia. *Journal of Tropical Ecology* 26:271-284.
- Pereira de Souza, S., S. Pacca, M. Turra de Avila, and J. L. B. Borges. 2010. Greenhouse gas emissions and energy balance of palm oil biofuel. *Renewable Energy* 35:2552-2561.
- Yamakura, T., A. Hagihara, and S. O. Sukardjo, H. 1986. Aboveground biomass of tropical rain forest stands in Indonesian Borneo. *Plant Ecology* 68:71-82.