

**Appendix 2.** Values used for the conversion of unit animal foods to weight and edible weight consumed by interviewed households (expressed in kg). Edible quantity was estimated from reference to the edible proportion of each species or group of species.

<i>Food items</i>	<i>Species (where applicable)</i>	<i>Weight per unit (Kg)</i>	<i>Edible weight (%)</i>	<i>Price per unit (Euro)</i>	<i>Price per kg (Euro)</i>	<i>References (weight and conversion)</i>
Fish		Variable	0.55	Varies	2,45	Torry res.stat. (1989) <sup>1</sup> , Holland et al. (1983) <sup>2</sup>
Chicken		1.300	0.65	3,5	2,65	Bender (1992) <sup>3</sup> , Chan et al. (1996) <sup>4</sup>
Duck		3.00	0.60	12,5	4,08	Albrechtsen et al.(2006) <sup>5</sup>
Domestic pig		48.30	0.60	Variable	4,08	Bender(1992) <sup>3</sup> , Chan et al. (1996) <sup>4</sup> , Bonfim (2002) <sup>6</sup>
Goat		25.00	0.60	50	3,27	Bender(1992) <sup>3</sup> , Chan et al. (1996) <sup>4</sup>
Sheep		20-30.00	0.60	60	4,08	Bender (1992) <sup>3</sup> , Chan et al. (1996) <sup>4</sup>
Egg		0.050	1.00	0,24	4,90	Holland et al. (1989) <sup>7</sup>
West African giant snail	<i>Archachatina marginata</i>	0.116	0.38	0,016	0,14	Ajayi et al. (1978) <sup>8</sup>
Obo snail	<i>Archachatina bicarinata</i>	0.116	0.38	na	na	Ajayi et al. (1978) <sup>8</sup>
Feral pig	<i>Sus domesticus</i>	50.00	0.60	Na	4,08	Bonfim (2002) <sup>6</sup>
Mona monkey	<i>Cercopithecus mona</i>	2.733	0.60	5 to 15	2,55	Albrechtsen et al.(2006) <sup>5</sup>
African civet	<i>Civectis civetta</i>	12.335	0.60	25	2,03	Albrechtsen et al.(2006) <sup>5</sup>
Fruit bat	<i>Eidolum helvum</i>	0.300	0.60	0,8	2,4	Albrechtsen et al.(2006) <sup>5</sup>
São Tomé green pigeon	<i>Treron sanctithomae</i>	0.230	0.50	1,2	5,32	Omojola et al.(2012) <sup>9</sup>
Bronze naped pigeon	<i>Columba malherbii</i>	0.165	0.50	1	6,06	Omojola et al.(2012) <sup>9</sup>
Maroon pigeon	<i>Columba thomensis</i>	0.400	0.50	1,5	3,75	Omojola et al.(2012) <sup>9</sup>

### References:

1. Torry Research Station, Aberdeen (1989). *Yield and nutritional value of the commercially more important fish species*.FAO Fisheries Technical Paper No. 309. FAO, Rome.
2. Holland, B., J. Brown, and D.H. Buss (1993). *Fish and Fish products*. Third supplement to the 5<sup>th</sup> edition of Mc Cance and Widdowson's *The Composition of Foods*. The Royal Society of Chemistry, Cambridge.

3. Bender, A. (1992). *Meat and meat products in human nutrition in developing countries*. Commissioned jointly by the Animal Production and Health Division and the Food Policy and Nutrition Division of FAO. Food and Nutrition Paper 53. FAO, Rome.
4. Chan, W., J. Brown, S.M. Church, and D.H. Buss (1996). *Meat Products and Dishes*. Sixth supplement to the 5<sup>th</sup> edition of Mc Cance and Widdowson's *The Composition of Foods*. The Royal Society of Chemistry, Cambridge.
5. Albrechtsen, L., J.E. Fa, B. Barry, and D.W. Macdonald (2006). Contrasts in availability and consumption of animal protein in Bioko Island, West Africa: the role of bushmeat. *Environmental Conservation*, 32: 340.
6. Bonfim, F. (2002) *Monografia sobre a Pecuária*. Estratégia Nacional e Plano de Acção da Biodiversidade, São Tomé.
7. Holland, B., I.D. Unwin, and D.H. Buss (1989). *Milk Products and Eggs*. Fourth supplement to the 5<sup>th</sup> edition of Mc Cance and Widdowson's *The Composition of Foods*. The Royal Society of Chemistry, Cambridge.
8. Ajayi, S. S., O. Tewet, C. Moriarty, and M.O. Awesu (1978). Observations on the biology and nutritive value of the African giant snail *Archachatina marginata*. *African Journal of Ecology*, 16(2): 85-95.
9. Omojola, A. B., M.A. Isa, M. Jibir, B.T. Ajewole, S. Garba, O.R. Kassim, A.B. Omotoso, O.A. Adeyemo, and S.B. Akinleye (2012). Carcass Characteristics and Meat Attributes of Pigeon (*Columbia Livia*) as Influenced by Strain and Sex. *Journal Animal Science Advances*, 2(5): 475-480.